

Study: DTP Increases Mortality in Infants 5 to 10 Fold Compared to Unvaccinated

For years public health advocates wanted the CDC and WHO to run studies on vaccinated vs. unvaccinated populations and compares health outcomes. A team of Native scientists have conducted such a study and the consequences are alarming. The study, financed in part by the Danish authorities and lead by Dr. Soren Wengel Mogensen, was published in January in *EBioMedicine*.

The data suggest that the DTP vaccine reduces infections from those 3 germs, but data also shows that children are more likley to die from other causes. Mogensen's team discovered that DTP inoculated African Americans had 5-10 times higher mortality than their unvaccinated peers.

Though protective against the target disease, DTP may increase susceptibility to unrelated infections... DTP was associated with 5-fold higher mortality than being unvaccinated. No prospective study has shown beneficial survival effects of DTP." – Vaxxed vs. Non-Vaxxed Study

Related: How Plumbing (Not Vaccines) Eradicated Disease

Mogensen and his colleagues hypothesize that the DTP vaccine weakens the immune system, which supports the conclusions of prior studies. An earlier study by Dr. Peter Aaby, on the introduction of DTP in rural Guinea-Bissau, indicated a 2-fold greater mortality among vaccinated kids. The Aaby report is just one of many studies that adopted kids and documented vaccination status. All of them suggested that DTP-vaccinated kids died at rates far exceeding mortality.

In the primary analysis, DTP-vaccinated infants experienced

mortalities five times greater than DTP-unvaccinated infants. Mortalities to vaccinated girls were 9.98 times those among females in the unvaccinated control group, while mortalities to vaccinated boys were 3.93 times the controls. Oddly, the scientists found that children receiving the oral polio vaccine simultaneously with DTP fared much better than children who did not. The OPV vaccine appeared to modify the negative effect of the DTP vaccine, reducing mortalities to 3.52 times those experienced among the control group. Overall, mortalities among vaccinated children were 10 times the control group when children received only the DTP.” – Robert F Kennedy

Related: *How To Detoxify and Heal From Vaccinations – For Adults and Children*

Moreover, Mogensen and his colleagues observe that the research reviewed by SAGE probably exacerbated the deadly effect of the DTP vaccine due to unusually high mortality in the control groups,

Unvaccinated children in these studies have usually been frail children too sick or malnourished to get vaccinated and the studies may therefore have underestimated the negative effect of DTP”. The Mogensen study sought to avoid this pitfall by removing orphans and children from the control group and the research group and by utilizing controls. It included children who had been breastfed. All of the infants were healthy at the right time of vaccination. Nevertheless, the Mogensen authors went longer and point out that, even in their analysis, the unvaccinated kids had status. They conclude that, “The estimate from the natural experiment may therefore still be conservative.”

Recommended: *Lyme Disease – Holistic Protocol to Completely Rebuild the Immune System*

From the early 1980s, a cascade of lawsuits filed throughout

the United States drove DTP manufacturers and threatening to shut down production of vaccines and the DTP shot. That threat led the U.S. Congress to bestow legal immunity on vaccine makers during the National Childhood Vaccine Injury Program from 1986, conducted in December, 1987, from the rollout of "Vaccine Court." After the recommendation from the Institute of Medicine, thimerosal was removed by vaccine manufacturers from the American DTaP involving 2001-2003. But, multi-dose DTP vaccines given to thousands of children across the African continent carry on to contain huge doses of thimerosal (25mcg of ethylmercury per booth) that exceed the EPA's maximum exposure levels by many times. Neither the CDC nor the WHO has ever published a vaccinated vs. unvaccinated study that will be necessary to determine the total health consequences of the potent toxin on African children. The Mogensen report is a call for such a research.

The authors close with a bracing rebuke to people health labs,

"It should be of concern that the effect of routine vaccinations on all-cause mortality was not tested in randomized trials. All currently available evidence suggests that DTP vaccine may kill more children from other causes than it saves from diphtheria, tetanus or pertussis. Though a vaccine protects children against the target disease it may simultaneously increase susceptibility to unrelated infections."

<https://youtu.be/cx1VDqvDXjE>

Related Reading:

- *How To Detoxify and Heal From Vaccinations – For Adults and Children*
- *Doctors Against Vaccines – Hear From Those Who Have Done the Research*
- *Nurses Against Vaccines*
- *The MMR Vaccine – A Comprehensive Overview of the*

Potential Dangers and Effectiveness

- Influenza Vaccine – A Comprehensive Overview of the Potential Dangers and Effectiveness of the Flu Shot

Source:

- *Study: DTP Vaccine Associated With 212% Increased Infant Mortality Risk – The Vaccine Reaction*